



BERES® - 8 SUPER HUMATE WITH FULVIC ACIDS AND MICROELEMENTS, UNIVERSAL CONCENTRATE WITH SULFUR

COMPOSITION:

	g/l	%
■ sulfur (SO ₃).....	300	25
■ humic acids.....	42	4
■ fulvic acids.....	18	1,7
■ amber acid.....	0,14	0,013
■ nitrogen (N).....	10,38	1

	г/л	%
■ potassium (K ₂ O).....	8,4	0,7
■ complex of macro- and microelements		
pH	8-8,5	
density	1,2 g/cm ³	

BERES® - 8 SUPER HUMATE WITH FULVIC ACIDS AND MICROELEMENTS, UNIVERSAL CONCENTRATE WITH SULFUR

organomineral fertilizer based on a sulfur compound that is easily accessible to plants, a highly active complex of fulvic and humic acids, amber acid, macro- and microelements. Anti-stress agent, growth and nutrition stimulator, adaptogen, immunomodulator.

Active components of the product:

- restore damaged cell elements,
- strengthen cell walls and limit the penetration of pathogens into plant tissues,
- improve the absorption of nitrogen compounds, which gives a visual increase in the intensity and uniformity of leaf color ("green effect"),
- prevent the formation of nitrates in agricultural products, ensuring their high environmental friendliness,
- Increase the content of proteins, vitamins, enzymes, oils and sugars.

Stimulates the growth of vegetative and generative mass. Increases the efficiency of photosynthesis and metabolic processes under normal and stressful conditions. Removes pesticide load from cultivated plants. Increases resistance to diseases, activating its own protective functions. Increases resistance to natural and abiotic stress. Helps increase productivity and improve product quality.

PREPARATION FORM:
liquid

PACKAGING:
canister
10 l, 5 l, 1 l

CONSUMPTION RATES:
0,5 - 1,5 l per 1 hectare of crops

Crop	Seed treatment	Application phases		
Spring and winter grains		tillering - beginning of stem elongation	flag leaf - ear formation	flowering - beginning of milky ripeness
Corn		appearance of 3-8 leaves	booting	heading of panicles
Buckwheat		first pair of true leaves - branching	budding	flowering, fruit formation
Peas, chickpeas, soybeans, lentils, beans		seedlings - leaves of the first tier	leaves of the second - fourth tier	budding - beginning of flowering, formation of pods
Rapeseed, mustard, winter cress spring and winter		formation of a leaf rosette - branching	stem growth - beginning of budding	budding - beginning of flowering, formation of pods
Flax, camelina		herringbone	budding, flowering	seed ripening
Sunflower		2-4 pairs of true leaves	6-8 pairs of true leaves	forming anantheode - beginning of flowering
Sugar beet and table beet		2-4 pairs of true leaves	4-8 pairs of true leaves - closing of crops in rows	closing of crops between rows
Potato		sprouting - plant height 10-15 cm	stem growth, budding	flowering - tuber formation
Solanaceae (tomatoes, peppers, eggplants)		appearance of 2-4 leaves	active growth - formation of set	filling of fruits - ripening
Cabbage		2-3 days after planting the seedlings	4-5 true leaves - beginning of glome setting	loaf formation
Carrot		sprouting - formation 1-2 true leaves	active leaf growth	root growth, root formation
Onion, garlic, radish		appearance of 2-3 leaves	active vegetative growth	beginning of formation - growth of root bulb
Fruit and berry		flower heads phase	before flowering	growth of fruit inception
Grapes		budding	after flowering	ripening of berries
Flower and decorative crops		sprouting - 2-3 leaves	appearance of 5-7 leaves	budding

HOW TO USE:

- foliar, root feeding, fertigation, drip irrigation - together with plant protection agents, or independent application.

